



Year: 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood in subalpine forests determined using dendrochronology and radiocarbon dating

Petrillo, Marta ; Cherubini, Paolo ; Fravolini, Giulia ; Ascher, Judith ; Schärer, Michael ; Synal, Hans-Arno ; Bertoldi, Daniela ; Camin, Federica ; Larcher, Roberto ; Egli, Markus

Abstract: Due to the large size and highly heterogeneous spatial distribution of deadwood, the time scales involved in the coarse woody debris (CWD) decay of *Picea abies* (L.) Karst. and *Larix decidua* Mill. in Alpine forests have been poorly investigated and are largely unknown. We investigated the CWD decay dynamics in an Alpine valley in Italy using the five-decay class system commonly employed for forest surveys, based on a macromorphological and visual assessment. For the decay classes 1 to 3, most of the dendrochronological samples were cross-dated to assess the time that had elapsed since tree death, but for decay classes 4 and 5 (poorly preserved tree rings) and some others not having enough tree rings, radiocarbon dating was used. In addition, density, cellulose and lignin data were measured for the dated CWD. The decay rate constants for spruce and larch were estimated on the basis of the density loss using a single negative exponential model. In the decay classes 1 to 3, the ages of the CWD were similar varying between 1 and 54 years for spruce and 3 and 40 years for larch with no significant differences between the classes; classes 1–3 are therefore not indicative for deadwood age. We found, however, distinct tree species-specific differences in decay classes 4 and 5, with larch CWD reaching an average age of 210 years in class 5 and spruce only 77 years. The mean CWD rate constants were 0.012 to 0.018 yr⁻¹ for spruce and 0.005 to 0.012 yr⁻¹ for larch. Cellulose and lignin time trends half-lives (using a multiple-exponential model) could be derived on the basis of the ages of the CWD. The half-lives for cellulose were 21 yr for spruce and 50 yr for larch. The half-life of lignin is considerably higher and may be more than 100 years in larch CWD.

DOI: <https://doi.org/10.5194/bgd-12-14797-2015>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-120425>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Petrillo, Marta; Cherubini, Paolo; Fravolini, Giulia; Ascher, Judith; Schärer, Michael; Synal, Hans-Arno; Bertoldi, Daniela; Camin, Federica; Larcher, Roberto; Egli, Markus (2015). Time since death and decay

rate constants of Norway spruce and European larch deadwood in subalpine forests determined using dendrochronology and radiocarbon dating. Biogeosciences Discussions, 12(17):14797-14832.
DOI: <https://doi.org/10.5194/bgd-12-14797-2015>



This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Time since death and decay rate constants of Norway spruce and European larch deadwood in subalpine forests determined using dendrochronology and radiocarbon dating

M. Petrillo^{1,2}, P. Cherubini², G. Fravolini⁴, J. Ascher^{5,6}, M. Schärer¹, H.-A. Synal³, D. Bertoldi⁷, F. Camin⁷, R. Larcher⁷, and M. Egli¹

¹Department of Geography, University of Zurich, 8057 Zurich, Switzerland

²WSL Swiss Federal Institute for Forest, Snow and Landscape Research, 8903 Birmensdorf, Switzerland

³Laboratory of Ion Beam Physics, ETH Zurich, 8093 Zurich, Switzerland

⁴Department of Bioscience and Territory, University of Molise, 86090 Pesche, Italy

⁵Department of Agrifood and Environmental Science, University of Florence, 50144 Florence, Italy

⁶Institute of Microbiology, University of Innsbruck, 6020 Innsbruck, Austria

14797

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁷IASMA, Fondazione Edmund Mach, 38010 San Michele all'Adige, Italy

Received: 13 July 2015 – Accepted: 20 August 2015 – Published: 8 September 2015

Correspondence to: M. Egli (markus.egli@geo.uzh.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Due to the large size and highly heterogeneous spatial distribution of deadwood, the time scales involved in the coarse woody debris (CWD) decay of *Picea abies* (L.) Karst. and *Larix decidua* Mill. in Alpine forests have been poorly investigated and are largely unknown. We investigated the CWD decay dynamics in an Alpine valley in Italy using the five-decay class system commonly employed for forest surveys, based on a macro-morphological and visual assessment. For the decay classes 1 to 3, most of the dendrochronological samples were cross-dated to assess the time that had elapsed since tree death, but for decay classes 4 and 5 (poorly preserved tree rings) and some others not having enough tree rings, radiocarbon dating was used. In addition, density, cellulose and lignin data were measured for the dated CWD. The decay rate constants for spruce and larch were estimated on the basis of the density loss using a single negative exponential model. In the decay classes 1 to 3, the ages of the CWD were similar varying between 1 and 54 years for spruce and 3 and 40 years for larch with no significant differences between the classes; classes 1–3 are therefore not indicative for deadwood age. We found, however, distinct tree species-specific differences in decay classes 4 and 5, with larch CWD reaching an average age of 210 years in class 5 and spruce only 77 years. The mean CWD rate constants were 0.012 to 0.018 yr^{-1} for spruce and 0.005 to 0.012 yr^{-1} for larch. Cellulose and lignin time trends half-lives (using a multiple-exponential model) could be derived on the basis of the ages of the CWD. The half-lives for cellulose were 21 yr for spruce and 50 yr for larch. The half-life of lignin is considerably higher and may be more than 100 years in larch CWD.

1 Introduction

The quantity and residence time of deadwood in Alpine forests are crucial in assessing the carbon cycle to ensure sustainable management of forests. Deadwood plays an important role in maintaining biodiversity in forest ecosystems (Müller and Bütler, 2010)

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as well as storing carbon (Di Cosmo et al., 2013), and contributing to nutrient cycle processes (Palviainen et al., 2010). The amount of deadwood varies greatly from managed to natural forests. In managed European Alpine forests, for example, the average stock of deadwood is estimated to be about $26 \text{ m}^3 \text{ ha}^{-1}$, while in old growth Alpine coniferous forests it can be up to $150\text{--}190 \text{ m}^3 \text{ ha}^{-1}$ (Barbati et al., 2014). Residence time for deadwood (e.g. Krüger et al., 2014) – from the moment the tree reaches the forest floor until it loses 95 % of the mass – can range from decades to several hundred years, depending on intrinsic and external factors. These factors include the dimensions of the log, the wood chemistry and the site conditions, in particular the mean annual temperature and soil moisture.

Until now, various different sampling designs have been used to determine the time since death to estimate the decay rate of deadwood. Long-term studies can provide reliable results (Müller-Using and Bartsch, 2009), but the slow decay dynamics of wood usually require a decadal observation period. Bond-Lamberty and Gower (2008) used the ratio of deadwood mass input into the pool of initial deadwood to estimate its decay rate based on a 7-years observation period. Such time sequences (chronosequence) offer ideal scenarios to study deadwood dynamics. If windthrow, fire regeneration and harvest events are known, the starting point in the timeline of the decay process can be specified. However, the exact year of such events is often uncertain which means precisely dating a tree's death is critical. Dendrochronology can be a helpful tool to determine the year of death, and the technique has been used in several studies to determine the time elapsed since tree-death (Campbell and Laroque, 2007; Lombardi et al., 2008, 2013). Other researchers have used radiocarbon dating to date the last recognizable ring of deadwood. For example, Kueppers et al. (2004) estimated the turnover time of lodgepole pine along a subalpine elevation gradient and Krüger et al. (2014) compared tree-ring cross-dating and radiocarbon dating, demonstrating that the two techniques produce comparable results. The decay rate can be estimated by relating the time-since-death to the density loss or mass loss of deadwood during a given time period (e.g. Busse, 1994; Melin et al., 2009). The decay rate is commonly expressed

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



through a decay constant k , which indicates the density loss or mass loss per year. This constant is derived from a decay model (Harmon, 1986), which can be most simply expressed by the equation

$$x_t = x_0 e^{-kt} \tag{1}$$

(single-negative-exponential model), where x_t is the density or mass of deadwood at a given time, and x_0 is the initial density or mass (Jenny et al., 1949; Olson, 1963). Other decay models have also been developed that take wood decomposition into account (reviewed by Mackensen et al., 2003). Some consider the different wood components, e.g. bark, sapwood, heartwood and chemical compounds, and combine them in multiple-exponential equations. Other models consider the time elapsed from the death of a standing tree to the moment when it falls and comes in contact with the forest floor (lag-time models). In several environments, e.g. on dry mountain slopes, the time lag between death and contact with the forest floor can last for almost the entire decay process (Kueppers et al., 2004). A few models take not only the losses due to heterotrophic respiration and leaching into account, but also losses due to fragmentation (Mackensen et al., 2003).

One of the most important components of deadwood is coarse woody debris (CWD). Because the spatial distribution of CWD is highly heterogeneous, only little quantitative data about its long-term decay dynamics are available for European Alpine forests. Decay models in Europe have, therefore, rarely been parameterised using empirically derived decay constants. In the field, the different stages of CWD decomposition are often described by so-called decay classes (as defined by Hunter, 1990) through a visual assessment of the wood status (Lombardi et al., 2013). In a previous study, Petrillo et al. (2015) demonstrated that the Hunter classification is particularly suitable for describing changes in the physical-chemical characteristics of European larch (*Larix decidua* Mill.) and Norway spruce (*Picea abies* (L.) Karst.) deadwood in alpine environments. The physical-chemical properties of deadwood changed distinctly during decay and correlated well with the 5 decay classes. Furthermore, no substantial differences

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

between spruce and larch decay patterns were found, although the wood chemistry of the living trees differed slightly between these two species. European larch and spruce are widespread in the Alps. Although C-stocks in soils are substantial (e.g. Johnston et al., 2004), CWD is a non-negligible C reservoir in subalpine forests (Sandström et al., 2007). Consequently, it is thus very important to know which time scales are involved in CWD decay. Jebrane et al. (2014) showed that Scots pine is more decay resistant than European larch, which suggests that the decay rate of pine is lower. Some species of larch are, however, considered economically valuable due to their hard, heavy and decay-resistant wood (Parker, 1993), which implies that residence time of larch CWD should be longer.

The aim of our work was to find out, (i) which time scales are involved in CWD decay of *P. abies* and *L. decidua* in the Alps and (ii) how these time scales correlate with the five-decay class system. We hypothesised that the CWD decay of these coniferous trees is relatively slow (due to, e.g. the nutrient availability for macro and micro-organisms being unfavourable) and comparable to soil organic matter compounds that are not easily degradable.

2 Materials and methods

2.1 Site description

The study area is located in the north-eastern Italian Alps, in Val di Sole and Val di Rabbi (Fig. 1; Table 1). The climate of the valleys ranges from temperate to alpine (above the timberline), the mean annual temperature from 8.2°C at the valley floor to about 0°C at 2400 m a.s.l., and the mean annual precipitation from approximately 800 to 1300 mm (Sboarina and Cescatti, 2004). The geological substrate is paragneiss debris in all sites. The soil units are Cambisols, Umbrisols and Podzols (WRB: IUSS working group, 2014). The soil properties at each site could be taken from a previous

study (Egli et al., 2006). The timberline is close to 2000–2200 m.a.s.l., with the forests dominated by Norway spruce and, at the highest altitudes, by European larch.

2.2 Sampling protocol

Norway spruce and European larch CWD was sampled at eight sites ranging in altitude from 1200 to 2000 m.a.s.l. In spring and summer 2013, wood cores from living trees and cross sections of CWD were taken from all sites. At each site, 5 or 6 living trees were sampled in two directions per each tree at 130 cm height (breast height) using an incremental corer (0.5 cm in diameter; Suunto, Finland). The wood cores were wrapped in paper and transported to the laboratory, where they were air dried, fixed onto a flat wooden support and sanded in order to obtain a smooth surface for tree-ring measurements. Before sampling, each CWD was first classified relative to the decay stage. The classification was done in situ using the five-class classification system of Hunter (1990) (Table 2), which is based on visual, geometric and tactile features and considers the presence/absence of twigs and bark, the shape of the log section, and the deadwood structure. Samples were taken randomly either using a manual saw or, in more advanced stages of decay, simply by hand. If necessary, they were wrapped up with tape to preserve their structure during transport to the laboratory, where they were air dried and sanded. For CWD in more advanced decay stages (decay class 4 and 5), a 25 cm × 30 cm bag was filled. The samples were then oven-dried at 50 °C, but not sanded. In total, 83 wood cores were taken from living trees, 29 from larch and 54 from spruce, while 40 cross sections were taken from deadwood (12 from larch and 28 from spruce).

2.3 Dendrochronological dating

At each site, the 10 or 12 wood cores taken from living trees were used to build a reference (master) ring-width chronology for each species. Tree rings were first counted and then measured using the LINTAB tree-ring-width measurement device (RINNTECH

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



e.K., Heidelberg, Germany), coupled together with a stereomicroscope (Leica, Germany). The two ring-width measurements from the same tree were first cross-checked and then incorporated into a single average master chronology for each species and for each site. The statistical software TSAP-winTM (Time Series Analysis Program, RIN-
NTECH e.K., Heidelberg, Germany) was used to calculate the *Gleichläufigkeit*, GLK (Kaennel and Schweingruber, 1995), i.e. the agreement between two ring-width series. The correlations among all the ring-width series of living trees and CWD were statistically assessed using the software COFECHA (Holmes et al., 1986).

The deadwood cross-sections were measured from the most external ring to the pith, along three or four different radial directions. The individual CWD series (i.e. floating chronologies) were matched to the master chronology of the corresponding species. We visually and statistically checked the deadwood series using the GLK to obtain the highest value with the master chronology and to date the year of death of the tree from which the deadwood originated.

2.4 Determining the cellulose and lignin

To obtain α -cellulose (Boettger et al., 2007), 10 mg of powdered wood were weighed in Teflon pockets for chemical and thermal treatments. Samples were first washed in a 5 % NaOH solution at 60 °C for two hours and then for an additional two hours with a 5 % NaOH solution, before finally being rinsed three times using boiling distilled water (see also Petrillo et al., 2015). The samples were then washed in a 7 % NaClO₂ solution at 60 °C for 30 h, changing the solution at least every 10 h and then rinsed three times with boiling distilled water. The pockets were dried in the oven at 50 °C and the cellulose content was determined as the difference between the initial weight and dried samples. The so-called Klason lignin (lignin insoluble in strong acid; Dence and Lin, 1992) was determined gravimetrically after a sequential extraction in which 0.2 g of each sample was washed three times with 5 mL of distilled water at 80 °C. After each washing, the samples were centrifuged for 10 min at 4500 rpm, dried in the oven at 80 °C and washed three times with 5 mL of ethanol. They were then centrifuged again

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(10 min at 4500 rpm) and the supernatant was discarded. After being dried at 60 °C in the oven, 60 mg of each sample were treated with 0.6 mL 72 % H₂SO₄ in a warm (30 °C) bath for one hour, and then, after adding 16.8 mL of distilled water, in an autoclave at 120 °C for one hour. Subsequently, the samples were filtered and the filtrate used to determine of the acid-soluble lignin. The insoluble lignin was dried in the oven at 105 °C and determined as the difference between the dry and initial weight.

2.5 Radiocarbon dating

The CWD of the decay classes 4 and 5 was too degraded to be dated through tree-ring analysis as their wood structure was too altered and the tree rings were no longer visible. In such cases, the outermost part of the CWD was sampled and ¹⁴C-dated (Fig. 2a and b). We selected a small fragment of 1 to 2 cm³ in volume from the outermost part assumed to have contained the last tree rings produced before the tree died (Fig. 2c and d). This small fragment was gently cleaned with a brush to remove any non-woody elements, such as particles of soil or vegetation like moss.

The organic samples were cleaned using an acid-alkali-acid (AAA) treatment. The samples were then heated under vacuum in quartz tubes with CuO (oxygen source) to remove any absorbed CO₂ in the CuO. The tubes were evacuated, sealed and heated in the oven at 900 °C to obtain CO₂. The CO₂ of the combusted sample was mixed with H₂ (1 : 2.5) and catalytically reduced over iron powder at 535 °C to elemental carbon (graphite). After reduction, the mixture was pressed into a target so that carbon ratios could be measured by Accelerator Mass Spectrometry (AMS) using the 0.2 MV radiocarbon dating facility (MICADAS) of the Laboratory of Ion Beam Physics at the Swiss Federal Institute of Technology of Zurich (ETHZ).

The calendar ages were obtained using the OxCal 4.2 calibration program (Bronk Ramsey, 2001, 2009) based on the IntCal 09 calibration curve (Reimer et al., 2013). Several samples (before 1950 AD) had a widely calibrated age range. For these samples, we used the age range with the highest probability of confining the time elapsed since death very strictly.

14805

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.6 Estimating decomposition rate constants on the basis of density loss

In a previous investigation (Petrillo et al., 2015), the density of the deadwood samples was measured. To estimate the decay constants, the average densities in class 1 and in class 5 (the earliest and latest decay stages) were used and the single-negative exponential model of Jenny et al. (1949) applied (see Eq. 1). Equation (1) was then solved for the decay constant k according to Eq. (2):

$$k = \frac{-\ln(x_t/x_0)}{t} \tag{2}$$

where x_t is the density of each deadwood sample at a given time (i.e. the estimated time elapsed since death), and x_0 the initial density (Table 3).

3 Results

3.1 Living chronologies

Two master chronologies for spruce and larch were obtained extending over 164 and 141 years, respectively (Fig. 3). The spruce chronology ranged from 1848 to 2012 AD (Fig. 3) and the larch chronology from 1871 to 2012 AD. The tree-ring widths of the same species correlated well among each other with a high GLK. The series inter-correlation coefficients obtained using COFECHA were 0.535 for spruce and 0.641 for larch (Pearson correlation, all series above 0.3281 were significant; 99 % confidence interval). Spruce and larch, however, had quite different growth patterns. The spruce chronology indicated that the trees grew homogeneously throughout the whole observation period, while larch seemed to be more sensitive to climate with marked high and low growth periods (positive and negative pointer years, Fig. 3). Furthermore, the negative and positive pointer years were not synchronous in the larch and spruce master chronologies. The growth pattern of some trees, however, differed considerably

from that of the master chronology, possibly due to the specific growth conditions of the individual trees, e.g. if their growth was very suppressed because of competition. Such outliers were excluded from the chronologies. The two master chronologies is described in Table 4.

3.2 Age of coarse woody debris (CWD)

Most of the samples of the decay classes 1–3 could be dendrochronologically dated, but those of decay classes 4 and 5 had to be radiocarbon dated because of the poorly preserved tree rings (Tables 5 and 6). In the first three decay classes, the CWD ages of spruce and larch seem to be in a similar range. The values vary from 1 to 54 years.

Interestingly, the average age of CWD does not seem to increase from class 1 to 3. The average age was around 10–20 years for all decay classes assuming a relatively fast decay. In decay classes 4 and 5, the average and maximum ages of CWD were usually higher for larch than for spruce. In decay class 4, spruce CWD has an average of about 42 years (median 43 years; Fig. 4) and larch CWD an average of 87 years (median 45 years). In decay class 5, the average age of spruce CWD increases to 77 years and the age of larch CWD to 210 years. This shows that larch wood, particularly in the decay classes 4 and 5, is much more resistant to rotting than spruce.

3.3 Relations between year since death, decay class and physical-chemical properties of deadwood

The physico-chemical data for the CWD are given in Petrillo et al. (2015) so that the density and the cellulose and lignin contents could be plotted as a function of the decay class and age of the CWD (Fig. 5). Since the relationship between the age of the CWD and physical-chemical characteristics was rather stochastic for the decay classes 1–3, they were grouped and their average was used for further analysis. The decrease in density and cellulose concentrations and the simultaneous increase in lignin definitely

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

proceed faster for the spruce CWD than for the larch CWD (Fig. 5). An exponential function best describes the trends in the cellulose and lignin concentrations with time.

Based on the density loss of spruce and larch CWD, the decay rate constants per year (yr^{-1}) were calculated for each dated sample. For spruce, we obtained an average value of $0.018 \text{ (yr}^{-1}\text{)}$ and for larch $0.012 \text{ (yr}^{-1}\text{)}$. The k values were non-normally distributed. Using the Kruskal–Wallis statistical test, we assessed the effects of the factors elevation, exposition, species and decay class on the k values. None of them significantly influenced the decay rate constant. Nonetheless, the range of k values on south-facing plots seem to be slightly higher than those on the north-facing plots, which suggests the decomposition rates are faster on south-exposed slopes (Fig. 6). In addition, the k values were estimated by comparing the CWD density with their age and by plotting an exponential regression curve (not shown). This approach resulted in lower k values: 0.012 yr^{-1} for spruce and 0.05 yr^{-1} for larch. The mean residence time and half-lives are summarised in Table 7.

4 Discussion

Although the five-decay class system is well suited to describe changes in the physical and chemical properties of deadwood (Lombardi et al., 2008), no real differences in the age of the CWD classes 1–3 could be found. The CWD in decay class 4 and 5 was, however, clearly older. This implies that the first three decay classes are not clearly related to deadwood age. Similarly, Lombardi et al. (2013) found no relationship between the age of CWD and the chemical properties of decay classes 1–3. The main explanation for this unexpected finding is that there is probably a time lag between the death of a standing tree and its contact with the soil (Kueppers et al., 2004; Zielonka, 2006; Lombardi et al., 2013). Standing dead trees, i.e. snags, can remain upright for several years and decay much more slowly than fallen dead trees (Yatskov et al., 2003). Angers et al. (2012), however, observed that the wood density in snags in boreal forests already decreases after a few years. Decay rates they calculated are

comparable to those in our study. The density loss in standing dead trees could be due to the activity of cerambycid larvae, while the activity of the wood decomposers, mainly fungi, was impeded in snags due to the lack of moisture. The discrepancy between the macromorphology of deadwood (and consequently decay class) and the age of dead wood seems therefore to be related to the individual tree death history. Shortly after tree death, In fact, the wood is rapidly colonized by fungi (Zielonka, 2006). The CWD in classes 4 and 5 showed a relation to deadwood age that seems to be species-specific since larch CWD is older than spruce in both classes. With respect to the CWD ages in our study, classes 1 to 3 appear to be a single group, while classes 4 and 5 are different. The oldest sample (larch CWD) was about 244 years old – a surprisingly old age for wood lying on the forest floor (i.e. not buried). Spruce CWD in decay classes 4 and 5 seem to be significantly younger than larch CWD. Few empirical assessments of time since the death of a tree have been made in Europe. Krüger et al. (2014) used both dendrochronology and radiocarbon dating to assess the time since death of Norway spruce in Bavarian forests. They estimated a total residence time of 61 to 62 years for this species. Our values are slightly lower. In Atlantic Canada, Campbell and Laroque (2007) found an age of 56 to 84 years (depending on the investigated sites) in the latest decay stage (decay class 5; Black spruce and Balsam fir). Lombardi et al. (2008) estimated stumps of beech and silver fir in decay class 3 to be 55 and 59 years which is close to our findings.

The decay rates reflect the determined ages of the CWD; and spruce therefore had a higher decay rate constant than larch. Consequently, decay rates are species specific due to, among others things, the initial differences in the physical-chemical properties of the wood of the living trees and in environmental factors. Larch has, for example, a higher density (cf. Fig. 5) and a lower nutrient content than spruce (Petrillo et al., 2015). Shorohova et al. (2014) also found that decay rates can strongly vary among tree species. The decay rate (i.e. 0.032 yr^{-1}) they found for spruce was slightly higher than that in our study (Fig. 6). The variability of the decay rates given in the literature may also arise from using different mathematical models or different methods to de-

BGD

Time since death and decay rate constants of Norway spruce and European larch deadwood

Title Page

Conclusions

Tables

◀

[Back](#)

Full Screen / Esc



Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

termine wood density or the age of the CWD. According to Hale and Pastor (1998), the decay rates of oak and maple logs (in a temperate forest) varied between 0.00 and 0.18 yr^{-1} (their dating of the logs, however, based on estimates). The decay rates of tree species in a Mediterranean area (Australia; Brown et al., 1996) varied in the range of 0.05 up to 0.22 yr^{-1} , while in a cool-continental climate (Alban and Pastor, 1993), decay rates were 0.042 and 0.055 for red and jack pine, respectively, 0.07 for spruce and 0.08 yr^{-1} for aspen. Fukusawa et al. (2014) estimated decay rates by using the annual input of CWD divided by the CWD accumulation, and obtained a value of 0.036 yr^{-1} . With the chronosequence approach, however, the rates were in the order of $0.020\text{--}0.023 \text{ yr}^{-1}$.

Using mass losses instead of density losses to estimate the decay rates may result in higher values, because the losses for fragmentation are added to the mineralisation losses (Yin, 1999). This might explain why our decay rate constants were lower than those in some other studies (Rock et al., 2008). Moreover, the decay rates are sensitive, at a regional scale, to climatic conditions such as temperature and precipitation (Shorohova et al., 2014), although the decay rates for a mean annual temperature of 0 to 10°C are, however, quite similar, and rates below 0.04 yr^{-1} are often reported (Mackensen et al., 2003). Soil temperature was found to be the main explanatory variable for differences in the decay rates of standard wood, such as aspen and pine (Risch et al., 2013). Although the data are too limited to draw clear conclusion, some of the differences in the decay rates we observed are likely to be due to environmental factors. On south-facing sites, for instance, we found that the decay rates were slightly, but not significantly, higher than those on north-facing sites (Fig. 6), which is comparable to Shorohova's et al. (2014) observations.

The concentrations of cellulose and lignin in the CWD are given as a function of time in Fig. 5. Due to the faster decomposition of cellulose, lignin is relatively enriched. Lignin, however, also decomposes with time. To unravel the decay behaviour of these compounds, a multiple-exponential model was applied (Means et al., 1985; Mackensen

et al., 2003), with the general form:

$$x = x_1 e^{-k_1 t} + x_2 e^{-k_2 t} \dots + x_n e^{-k_n t} \quad (3)$$

where x_t is the density or mass of deadwood at a given time and $x_{1...n}$ are partitioned parameters. The portioning of cellulose and lignin is solved graphically using their mass per unit volume over time, and fitting them to an exponential regression curve. From this, the half-life of cellulose or lignin in the CWD could be calculated:

$$t_{1/2} = \frac{\ln(1/2)}{-k} \quad (4)$$

where $t_{1/2}$ is the half-life and k is the decay constant (obtained from the exponential regression curve). Using the k values in Fig. 7, the following half-lives were obtained:

1. for cellulose: 21 years (spruce) and 50 years (larch)
2. for lignin: 91 years (spruce) and 481 years (larch)

Means et al. (1985) were able to derive k values for cellulose values of 0.0109 to 0.0117 yr^{-1} for Douglas-fir logs (in a cool to temperate climate), although age determination (or estimation) was done differently. This would give rise to half-lives in the range of 59 to 64 years. With k values in the range of 0.0039 to 0.0045 yr^{-1} (Means et al., 1985), the half-life of lignin would be in the range of 154–178 years. In this specific case, the overall decay rates were between 0.006 and 0.0073 yr^{-1} . Although cellulose is relatively easily degradable by (micro)organisms, it may persist astonishingly long in larch trees (several decades). Lignin may have a half-life of more than hundred years. These half-lives may be shorter if the decay is related to mass losses and not to density.

How do such data now compare to organic matter in soils? In decay classes 4 and 5, the CWD starts to become more and more part of the soil. The introduced organic matter into soils can be either further degraded or stabilised to a certain extent. The persistence of organic matter in soils (SOM) is largely due to complex interactions between

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the organic matter and its environment, such as the interdependence of compound chemistry, reactive mineral surfaces, climate, water availability, soil acidity, soil redox state and the presence of potential degraders in the immediate micro-environment (Schmidt et al., 2011). Most organic components in soils have a mean residence time of just a few years up to about 50–60 years. Fire-derived organic matter may have a residence time of decades to as much as 300 years (Schmidt et al., 2011). The measured mean ages of the CWD in decay class 4 or 5, the CWD residence time and the decay rates of cellulose and lignin are very similar of those of soil organic matter compounds (and even fire-derived SOM). The rate of CWD decay processes does not apparently differ very greatly from the degradation of organic matter in soils.

5 Conclusions

The first 3 decay classes do not seem to reflect the age of the CWD, but they are relevant for the description of its decay stage. Taking these classes as one group and relating them to the decay classes 4 and 5, a time trend with increasing decay stage can then be detected. This time trend also closely correlates to the wood density, and the cellulose and lignin content. The oldest CWD age of a larch tree reached the considerable age of 244 years. The ages of the CWD in decay classes 4 and 5 even correspond to relatively “stable” SOM (e.g. fire-derived organic matter). Consequently, the turn-over time of CWD organic matter is in a similar range to that of SOM. The decay rate constant for spruce was 0.012 to 0.018 (yr^{-1}) and for larch 0.005 to 0.012 (yr^{-1}), depending on the approach. The rates seemed to be slightly higher on south-facing sites (although this was not statistically significant). Using the dating approach (dendrochronology and ^{14}C -dating), the behaviour of cellulose and lignin as a function of time could be assessed. Our findings demonstrate that lignin in larch may persist particularly long, with a mean residence time of > 100 years.

More empirical data is, however, needed to ascertain our findings. The preparation and precise dating of CWD is time-consuming and cost intensive. Since CWD repre-

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sents an important forest carbon pool, improving the informative potential of the decay classes (including the dating of the CWD) would contribute to sustainable forest management and make carbon accounting easier.

Acknowledgements. This study is part of the DecAlp DACH project no. 205321L_141186. J. Ascher has been funded by the Fonds zur Förderung der wissenschaftlichen Forschung (FWF) Austria (Project I989-B16). We are indebted to Fabio Angeli of the “Ufficio distrettuale forestale di Malé” and his team of foresters for their support in the field. We would also like to thank Leonora Di Gesualdo for her help in the sampling wood cores and Michelle Kovacic for preparing samples for radiocarbon dating. We are grateful to Silvia Dingwall for the English corrections.

References

Alban, D. H. and Pastor, J.: Decomposition of aspen, spruce, and pine boles on two sites in Minnesota, Can. J. Forest Res., 23, 1744–1749, doi:10.1139/x93-220, 1993.

Angers, V. A., Drapeau, P., and Bergeron, Y.: Mineralization rates and factors influencing snag decay in four North American boreal tree species, Can. J. Forest Res., 42, 157–166, doi:10.1139/X11-167, 2012.

Barbati, A., Marchetti, M., Chirici, G., and Corona, P.: European forest types and forest Europe SFM indicators: tools for monitoring progress on forest biodiversity conservation, Forest Ecol. Manag., 321, 145–157, doi:10.1016/j.foreco.2013.07.004, 2014.

Bond-Lamberty, B. and Gower, S. T.: Decomposition and fragmentation of coarse woody debris: re-visiting a boreal black spruce chronosequence, Ecosystems, 11, 831–840, doi:10.1007/s10021-008-9163-y, 2008.

Boettger, T., Haupt, M., Knoller, K., Weise, S. M., Waterhouse, J. S., Rinne, K. T., Loader, N. J., Sonninen, E., Jungner, H., Masson-Delmotte, V., Stievenard, M., Guillemain, M. T., Pierre, M., Pazdur, A., Leuenberger, M., Filot, M., Saurer, M., Reynolds, C. E., Helle, G., and Schleser, G. H.: Wood cellulose preparation method and mass spectrometric analysis of delta C-13, delta O-18, and nonexchangeable delta H-2 values in cellulose, sugar and starch: an interlaboratory comparison, Anal. Chem., 79, 4603–4612, doi:10.1021/ac0700023, 2007.

Bronk Ramsey, C.: Bayesian analysis of radiocarbon dates, Radiocarbon 51, 337–360, 2009.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Brown, S., Mo, J., McPherson, J. K., and Bell, D. T.: Decomposition of woody debris in Western Australian forests, *Can. J. Forest Res.*, 26, 954–966, doi:10.1139/x26-105, 1996.
- Campbell, L. J. and Laroque, C. P.: Decay progression and classification in two old-growth forests in Atlantic Canada, *Forest Ecol. Manag.*, 238, 293–301, doi:10.1016/j.foreco.2006.10.027, 2007.
- Dence, C. W. and Lin, S. Y.: Introduction, in: *Methods in Lignin Chemistry*, edited by: Lin, S. Y. and Dence, C. W., Springer, Heidelberg 1–19, 1992.
- Di Cosmo, L., Gasparini, P., Paletto, A., and Nocetti, M.: Deadwood basic density values for national-level carbon stock estimates in Italy, *Forest Ecol. Manag.*, 295, 51–58, doi:10.1016/j.foreco.2013.01.010, 2013.
- Egli, M., Mirabella, A., Sartori, G., Zanelli, R., and Bischof, S.: Effect of north and south exposure on weathering rates and clay mineral formation in Alpine soils, *Catena*, 67, 155–174, doi:10.1016/j.catena.2006.02.010, 2006.
- Hale, C. M. and Pastor, J.: Nitrogen content, decay rates, and decompositional dynamics of hollow versus solid hardwood logs in hardwood forests of Minnesota, U.S.A., *Can. J. Forest Res.*, 28, 1276–1285, doi:10.1139/cjfr-28-9-1276, 1998.
- Harmon, M. E., Franklin, J. F., Swanson, F. J., Sollins, P., Gregory, S. V., Lattin, J. D., Anderson, N. H., Cline, S. P., Aumen, N. G., Sedell, J. R., Lienkaemper, G. W., Cromack, K., and Cummins, K. W.: Ecology of coarse woody debris in temperate ecosystems, *Adv. Ecol. Res.*, 15, 133–302, doi:10.1016/S0065-2504(08)60121-X, 1986.
- Holmes, R. L., Adams, R. K., and Fritts, H. C.: Tree-ring chronologies of western North America: California, eastern Oregon and northern Greate Basin, with procedures used in chronology development work, including users manuals for computer programs COFECHA and ARSTAN, in: *Chronology series VI*, Laboratory of Tree-Ring Research, University of Arizona, Tucson, 182 pp., 1986.
- Hunter, M. L.: *Wildlife, forests and forestry: principles of managing forests for biological diversity*. Prentice Hall, Englewood Cliffs (NJ), USA, 370 pp., 1990.
- IPCC: *Good Practice Guidance for Land Use, Land Use Change and Forestry*, Institute for Global Environmental Strategies, Hayama, 2003.
- IUSS Working Group WRB: *World Reference Base for Soil Resources 2014*. International Soil Classification System for Naming Soils and Creating Legends for Soil Maps, World Soil Resources Reports No. 106, FAO, Rome, 2014.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Jebrane, M., Pockrandt, M., and Terziev, N.: Natural durability of selected larch and Scots pine heartwoods in laboratory and field tests, *Int. Biodeter. Biodegr.*, 91, 88–96, doi:10.1016/j.ibiod.2014.03.018, 2014.
- Jenny, H., Gessel, S. P., and Bingham, F. T.: Comparative study of decomposition of organic matter in temperate and tropical regions, *Soil Sci.*, 68, 419–432, doi:10.1097/00010694-200606001-00017, 1949.
- Kaennel, M. and Schweingruber, F. H.: Multilingual glossary of dendrochronology. Terms and definitions in English, German, French, Spanish, Italian, Portuguese and Russian, Swiss Federal Institute for Snow, Forest and Landscape Research, Haupt, Bern, Switzerland, 1995.
- Kueppers, L. M., Southon, J., Baer, P., and Harte, J.: Dead wood biomass and turnover time, measured by radiocarbon, along a subalpine elevation gradient, *Oecologia*, 141, 641–651, doi:10.1007/s00442-004-1689-x, 2004.
- Krüger, I., Muhr, J., Hartl-Meier, C., Schulz, C., and Borken, W.: Age determination of coarse woody debris with radiocarbon analysis and dendrochronological cross-dating, *Eur. J. Forest Res.*, 133, 931–939, doi:10.1007/s10342-014-0810-x, 2014.
- Lombardi, F., Cherubini, P., Lasserre, B., Tognetti, R., and Marchetti, M.: Tree rings used to assess time since death of deadwood of different decay classes in beech and silver fir forests in the central Apennines (Molise, Italy), *Can. J. Forest Res.*, 38, 821–833, doi:10.1139/X07-195, 2008.
- Lombardi, F., Cherubini, P., Tognetti, R., Cocozza, C., Lasserre, B., and Marchetti, M.: Investigating biochemical processes to assess deadwood decay of beech and silver fir in Mediterranean mountain forests, *Ann. For. Sci.*, 70, 101–111, doi:10.1007/s13595-012-0230-3, 2013.
- Mackensen, J., Bauhus, J., and Webber, E.: Decomposition rates of coarse woody debris – a review with particular emphasis on Australian tree species, *Aust. J. Bot.*, 51, 27–37, doi:10.1071/BT02014, 2003.
- Means, J. E., Cromack, K., MacMillan, P. C.: Comparison of decomposition models using wood density of Douglas-fir logs, *Can. J. Forest Res.*, 15, 1092–1098, doi:10.1139/x85-178, 1985.
- Moroni, T. M., Hagemann, U., and Beilman, D. W.: Dead wood is buried and preserved in a Labrador boreal forest, *Ecosystems*, 13, 452–458, doi:10.1007/s10021-010-9331-8, 2010.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Müller, J. and Bütler, R.: A review of habitat thresholds for dead wood: a baseline for management recommendations in European forests, *Eur. J. Forest Res.*, 129, 981–992, doi:10.1007/s10342-010-0400-5, 2010.
- Müller-Using, S. and Bartsch, N.: Decay dynamic of coarse and fine woody debris of a beech (*Fagus sylvatica* L.) forest in Central Germany, *Eur. J. Forest Res.*, 128, 287–296, doi:10.1007/s10342-009-0264-8, 2009.
- Olson, J. S.: Energy storage and the balance of producers and decomposers in ecological systems, *Ecology*, 44, 322–331, doi:10.2307/1932179, 1963.
- Parker, W. H.: Larix, in: *Flora of North America North of Mexico*, Vol. 2, edited by: Flora of North America Editorial Committee, Oxford University Press, 366–368, 1993.
- Palviainen, M., Finén, L., Laiho, R., Shorohova, E., Kapitsa, E., and Vanha-Majamaa, I.: Phosphorus and base cation accumulation and release patterns in decomposing Scots pine, Norway spruce and silver birch stumps, *Forest Ecol. Manag.*, 260, 1478–1489, doi:10.1016/j.foreco.2010.07.046, 2010.
- Petrillo, M., Cherubini, P., Sartori, G., Abiven, S., Ascher, J., Bertoldi, D., Camin, F., Barbero, A., Larcher, R., and Egli, M.: Decomposition of Norway spruce and European larch coarse woody debris (CWD) in relation to different elevation and exposition in an Alpine setting, *iForest*, doi:10.3832/for1591-008, 2015.
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hafflidason, H., Hajdas, I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Nui, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A., Turney, C., and van der Plicht, J.: IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP, *Radiocarbon*, 55, 1869–1887, doi:10.2458/azu_js_rc.55.16947, 2013.
- Risch, C. A., Jurgensen, F. M., Page-Dumroese, D. S., and Schütz, M.: Initial turnover rates of two standard wood substrates following land-use change in subalpine ecosystems in the Swiss Alps, *Can. J. Forest Res.*, 43, 901–910, doi:10.1139/cjfr-2013-0109, 2013.
- Rock, J., Badeck, F., and Harmon, M. E.: Estimating decomposition rate constants for European tree species from literature sources, *Eur. J. Forest Res.*, 127, 301–313, doi:10.1007/s10342-008-0206-x, 2008.
- Sboarina, C. and Cescatti, A.: Il clima del Trentino – Distribuzione spaziale delle principali variabili climatiche, Report 33, Centro di Ecologia Alpina Monte Bondone, Trento, Italy, 2004.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49–56, doi:10.1038/nature10386, 2011.
- 5 Shorohova, E. and Kapitsa, E.: Influence of the substrate and ecosystem attributes on the decomposition rates of coarse woody debris in European boreal forests, *Forest Ecol. Manag.*, 315, 173–184, doi:10.1016/j.foreco.2013.12.025, 2014.
- WRB, IUSS Working Group: World Reference Base for Soil Resources, 2nd edn., World Soil Resources Reports No. 103, First update, FAO, Rome, 2007.
- 10 Yatskov, M., Harmon, M. E., and Krankina, O. N.: A chronosequence of wood decomposition in the boreal forests of Russia, *Can. J. Forest Res.*, 33, 1211–1226, doi:10.1139/X03-033, 2003.
- Yin, X.: The decay of forest woody debris: numerical modelling and implications based on some 300 data cases from North America, *Oecologia*, 121, 81–98, doi:10.1007/s004420050909, 15 1999.
- Zielonka, T.: When does dead wood turn into a substrate for spruce replacement?, *J. Veg. Sci.*, 17, 739–746, doi:10.1111/j.1654-1103.2006.tb02497.x, 2006.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Characteristics of the study sites.

Plot ID	Elevation (m a.s.l.)	Aspect (° N)	Slope (°)	MAP* (mm yr ⁻¹)	Parent material	Dominating tree species	Land use	Soil classification (WRB) (Egli et al., 2006)
<i>North-facing sites</i>								
N01	1180	340	31	950	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Episkeletic Cambisol (Dystric)
N02	1390	0	28	1000	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Episkeletic Cambisol (Dystric)
N03	1620	0	29	1060	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Chromi-Endoskeletal Cambisol (Dystric)
N04	1930	20	12	1180	Paragneiss debris, moraine material	<i>Larix decidua</i>	Originally used as pasture	Episkeletic Podzol
<i>South-facing sites</i>								
S06	1185	160	31	950	Paragneiss debris	<i>Picea abies</i>	Ex-coppice, natural forest (ecological forestry)	Episkeleti-Endoleptic Cambisol (Chromi- Dystric)
S07	1400	145	33	1000	Paragneiss debris	<i>Larix decidua</i>	Natural forest (ecological forestry)	Dystric-Endoskeletal Cambisol
S08	1660	210	33	1060	Paragneiss debris	<i>Picea abies</i>	Natural forest (ecological forestry)	Skeletal Umbrisol
S09	1995	160	25	1180	Paragneiss debris	<i>Larix decidua</i>	Ex pasture, natural forest	Skeletal Umbrisol

* MAP = mean annual precipitation (Sboarina and Cescatti, 2004).

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I ◀](#)
[▶ I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 2. The five decay-class system of log decomposition (according to Hunter, 1990).

Log features	Decay classes 1	2	3	4	5
Bark	Intact	Partially absent	Absent	Absent	Absent
Twigs	Present	Partially absent or absent	Absent	Absent	Absent
Shape of radial section	Round	Round	Round	Oval	Very oval
Colour	Original	Original	Faded in the external part	Reddish brown or faded	Reddish or faded
Texture of wood	Intact	Intact	Soft outer layer, intact inner part	Small pieces, soft	Powdery or fibrous, very soft
Contact with soil	Log elevated on what remains of branches	Log in contact with soil	Log in contact with soil	Log in contact with soil	Log in contact with soil and partially buried

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Wood density values for Norway spruce and European larch given by the IPCC (2003) and in previous investigations (Petrillo et al., 2015).

	Wood density (g cm^{-3})	
	IPCC (2003)	Petrillo et al. (2015)
<i>Larix decidua</i> L.	0.46	0.59
<i>Picea abies</i> L. Karst	0.40	0.45

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Table 4. Description of the two master chronologies. The inter-series correlation is obtained with the software COFECHA.

Species	Number of dated series	Series inter-correlation	Length (Years AD)
Norway spruce	53	0.53	1848–2012
European larch	28	0.64	1871–2012

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 5. Ages of Norway spruce and European larch CWD in class 1–3 obtained mostly from dendrochronological measurements and a few (*) from ^{14}C -dating (cf. Appendix Table A1).

Sample ID	Plot	Tree species	Decay class	Year of death	CWD age
233	N03	Norway spruce	1	2009	4
4a	S08	Norway spruce	1	1992	21
213a	N01	Norway spruce	1	1988	25
217a	N02	Norway spruce	1	1969	44
99Aa	N03	Norway spruce	2	2006	7
219a	N03	Norway spruce	2	2004	9
232a	N03	Norway spruce	2	2004	9
80Ax	S08	Norway spruce	2	2003	10
201a	N02	Norway spruce	2	1996	17
200a	N03	Norway spruce	2	1993	20
216a	N01	Norway spruce	2	1970	43
204a	N02	Norway spruce	2	1959	54
218a	N03	Norway spruce	3	2012	1
220a	N03	Norway spruce	3	2005	8
226a	N03	Norway spruce	3	2005	8
15a	N01	Norway spruce	3	1979	34
191b	N02	Norway spruce	3	1970	43
236a	S09	European larch	1	2010	3
230a	N04	European larch	1	1973	40
L_10_c1_1*	S07	European larch	1	2007	6
91Bx	S07	European larch	2	2010	3
52Bv	S09	European larch	2	2000	13
S07_dc2_92*	S07	European larch	2	2003	10
S07_dc3_96*	S07	European larch	3	2004	9
S09_cl3_46*	S09	European larch	3	1973	40
S09_cl3_48*	S09	European larch	3	1968	45

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 6. Radiocarbon data of the deadwood samples (decay classes 4 and 5) as a function of site and tree species.

UZH number	ETH number	Sample code	Site	Tree species	Decay class	^{14}C -age	$\pm 1\sigma$	$\delta^{13}\text{C}$	$\pm \delta^{13}\text{C}$	cal AD	Average age ¹	cal AD ²	Average age ¹
								‰	‰	2 σ	years	years (probability)	years
UZ 6210	ETH-56612	37A	N4	European larch	4	-2931	29	-23.6	1.0	1962–1974	45		
UZ 6211	ETH-56613	34A	N4	European larch	4	170	31	-24.4	1.0	1658–1950	209	1720–1819 (50 %)	244
UZ 6213	ETH-56615	50A	S9	European larch	4	-659	31	-26.5	1.0	1957–2003	33		
UZ 6219	ETH-56853	95A	S7	European larch	4	-860	25	-25.5	1.0	1957–1998	36		
UZ 6227	ETH-56861	209	S7	European larch	4	-2545	25	-25.3	1.0	1962–1976	44		
UZ 6228	ETH-56862	214	S7	European larch	4	60	25	-23.9	1.0	1695–1919	206	1867–1919 (53.3 %)	120
UZ 6212	ETH-56614	45B	S9	European larch	5	183	31	-27.0	1.0	1650–1950	213	1726–1815 (51.5 %)	243
UZ 6224	ETH-56858	202	S9	European larch	5	140	25	-28.9	1.0	1669–1944	207	1798–1944 (52.6 %)	142
UZ 6264	ETH-60747	NO4_cl5_33	N4	European larch	5	185	25	-29.4	1.0	1656–1950	220	1728–1810 (54.7 %)	244
UZ 6214	ETH-56616	69A	S6	Norway spruce	4	-1331	30	-31.3	1.0	1958–1989	40		
UZ 6215	ETH-56849	72A	S6	Norway spruce	4	-2120	25	-23.6	1.0	1961–1980	43		
UZ 6216	ETH-56850	84A	S8	Norway spruce	4	-4080	25	-27	1.0	1966–1967	47		
UZ 6220	ETH-56854	97B	N2	Norway spruce	4	-3720	25	-31.4	1.0	1967–1968	46		
UZ 6221	ETH-56855	98A	N3	Norway spruce	4	-290	25	-19.7	1.0	1955–2009	31		
UZ 6226	ETH-56860	206	N3	Norway spruce	4	-2150	25	-23.9	1.0	1962–1979	43		
UZ 6217	ETH-56851	87A	S8	Norway spruce	5	130	25	-26.1	1.0	1677–1940	205	1800–1940 (59.5 %)	143
UZ 6218	ETH-56852	89A	S8	Norway spruce	5	-615	25	-24.6	1.0	1956–2004	33		
UZ 6222	ETH-56856	106A	N1	Norway spruce	5	-1665	25	-25.2	1.0	1959–1984	42		
UZ 6223	ETH-56857	197	N1	Norway spruce	5	-4595	25	-24	1.0	1963–1965	49		
UZ 6225	ETH-56859	205	N2	Norway spruce	5	10	25	-22.9	1.0	1699–1916	206	1879–1916 (82.2 %)	116

¹ Calculated as the mean value between the maximum and minimum age (2σ). For this range of years (2σ), associated probabilities are summed to 95.4 %.

² For samples dated in the period before the bomb peak and giving a wide calibrated age range, the age is also calculated for the most important calibrated time-range associated to the highest, corresponding probability (in brackets).

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 7. CWD decay parameters based on Eq. (1) (a) and on the regression approach (b).

	Decay constant k (yr^{-1})	Residence time (yr)	Half-life (yr)
a)			
Norway spruce	0.018	56	39
European larch	0.012	83	58
b)			
Norway spruce	0.012	84	58
European larch	0.005	222	154

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Table A1. Radiocarbon data of the deadwood samples of the decay classes 1–3.

UZH number	ETH number	Sample code	Site	Tree species	Decay class	C14 age	$\pm 1\sigma$	$\delta^{13}\text{C}$ ‰	$\pm \delta^{13}\text{C}$ ‰	Cal AD $\pm 1\sigma$	Average age years
UZ-6258	ETH-60741	L_10_c1_1	S7	European Larch	1	–435	25	–25.7	1	2006–2009	6
UZ-6260	ETH-60743	S07_dc2_92	S7	European Larch	2	–590	25	–26.9	1	2002–2004	10
UZ-6261	ETH-60744	S07_dc3_96	S7	European Larch	3	–545	25	–26.4	1	2003–2005	9
UZ-6262	ETH-60745	S09_cl3_46	S9	European Larch	3	–2865	25	–29.3	1	1973–1974	40
UZ-6263	ETH-60746	S09_cl3_48	S9	European Larch	3	–2775	25	–23.8	1	1962–1974	45

* Calculated as the mean value between the maximum and minimum age (1σ). For this range of years (1σ), associated probabilities summed to 68.2%.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.




Figure 1. Location of the study area with the major vegetation units and investigation sites. Data source: Museo delle Scienze (Trento), CORINE Landcover (Joint Research Center of the European Union) and scilands GmbH.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.




Figure 2. Cross section of **(a)** spruce deadwood in the field (site N03) and **(b)** larch deadwood (site S07). Examples **(c)** and **(d)** of deadwood fragments classified as decay class 4 dated using radiocarbon (outermost part of the wood piece).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.




Figure 3. Master chronologies for spruce (continuous line) and larch (dashed line) to cross-date the deadwood.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Figure 4. Box plots of the larch and spruce deadwood age as a function of decay class.

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 5. Relation between the age of spruce and larch CWD and density **(a)**, cellulose % **(b)** and lignin % **(c)**. The decay classes 1–3 were grouped together due to their similar age (Fig. 4).




Figure 6. Calculated decay rate constants (k) as a function of tree species and site exposure.

BGD

12, 14797–14832, 2015

Time since death and decay rate constants of Norway spruce and European larch deadwood

M. Petrillo et al.




Figure 7. Empirically determined exponential regression curves (principle of multiple-exponential model) for partitioning the decay behaviour of cellulose and lignin.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

